

A Numerically Verifiable Proof for M-theory Compactifications*

Michael Douglas¹, Daniel Platt², Yidi Qi³

¹Harvard University, Stony Brook University ²King's College London ³Northeastern University

Motivation

Y Calabi-Yau 3-fold (real dimension 6) with involution σ

Theorem (Joyce-Karigiannis [1]):

If $fix(\sigma)$ admits a **nowhere vanishing harmonic vector field**, then $(Y \times S^1)/\langle \sigma \rangle$ admits a metric with holonomy G_2 .

These manifolds are needed in M-theory (a flavour of string theory)

Problem: **only conjectural examples** are known

Goal: prove one of the conjectural examples works!

Background

Calabi-Yau manifold Y in $\mathbb{C}P^5$ be (a small perturbation of)

$$Y_0 := \{ \vec{z} \in \mathbb{C}P^5 : p(\vec{z}) = 0 \text{ and } q(\vec{z}) = 0 \}$$

where p is degree 2 polynomial, q is degree 4 polynomial.


$\sigma : Y \rightarrow Y$ the conjugation map, then "real locus" is

$$Y_{\mathbb{R}} := fix(\sigma) = \{ \vec{z} = (z_0 : \dots : z_6) \in Y : z_0, \dots, z_6 \in \mathbb{R} \}$$

Y has two metrics:

g_{CY}	g_{approx} (see [2])
Calabi-Yau metric, has $Ricci(g_{CY}) = 0$, no closed form expression known	An approximation of g_{CY} , has $Ricci(g_{approx}) \neq 0$ but small , very explicit formulae given

Y in orange,
 $Y_{\mathbb{R}}$ in green




Question

Does $Y_{\mathbb{R}}$ admit a **harmonic vector field with no zeros**?

$$V \in \Gamma(TY_{\mathbb{R}}), \quad \Delta_{CY}(V) = 0 \quad ?$$

Problem: Δ_{CY} depends on the non-explicit metric g_{CY}



A vector field \tilde{V} on $Y_{\mathbb{R}}$ which is harmonic with respect to g_{approx} . It can be computed explicitly and is nowhere 0, with a good bound from below: $|\tilde{V}| > \epsilon$.

Our approach

1. Find harmonic (unit norm) vector field $\tilde{V} \in \Gamma(TY_{\mathbb{R}})$ w.r.t. g_{approx} with $|\tilde{V}| > \epsilon$ **everywhere**.

2. ("Stability estimate") Find $C > 0$ with

$$\|g_{CY} - g_{approx}\| \leq C \cdot \|Ricci(g_{approx})\|$$

3. Classical fact (Hodge theory): ex. $V \in \Gamma(TY_{\mathbb{R}})$ that is g_{CY} -harmonic in same cohomology class as \tilde{V}

4. Find $D > 0$ with

$$\|V - \tilde{V}\| \leq D \cdot \|g_{CY} - g_{approx}\|$$


5. Conclusion: if

$$\|Ric(g_{approx})\| < \epsilon \cdot \frac{1}{C} \cdot \frac{1}{D}$$

then

$$|V| > |\tilde{V}| - |V - \tilde{V}| > \epsilon - \epsilon \cdot \frac{1}{C} \cdot \frac{1}{D} \cdot C \cdot D = 0$$

V has no zeros!



The perturbed vector field V on $Y_{\mathbb{R}}$ which is harmonic with respect to g_{approx} . If $\|Ricci(g_{approx})\|$ is very small, then the metrics g_{CY} and g_{approx} are so close to each other that harmonic vector fields of the two metrics are almost the same. Because \tilde{V} was nowhere 0, the new V is nowhere 0, even though the bound from below may be worse.

References

- [1] Dominic Joyce and Spiro Karigiannis. A new construction of compact torsion-free G_2 -manifolds by gluing families of Eguchi-Hanson spaces. *J. Differential Geom.*, 117(2):255–343, 2021.
- [2] Michael Douglas, Subramanian Lakshminarasimhan, and Yidi Qi. Numerical calabi-yau metrics from holomorphic networks. In *Mathematical and Scientific Machine Learning*, pages 223–252. PMLR, 2022.